

## 第七章参考文献

1. 姚泰主编. 生理学（七年制规划教材）. 北京: 人民卫生出版社, 2001
2. 姚泰主编. 人体生理学. 第3版. 北京: 人民卫生出版社, 2001
3. 本郷利憲, 廣重力主編. 標準生理學. 東京: 醫學書院, 2000
4. Guyton AC, Hall JE. Textbook of Medical Physiology. 11th edition, Philadelphia: Saunders, 2006
5. Anuurad E, Shiwaku K, Nogi A, et al. The new BMI criteria for asians by the regional office for the western pacific region of WHO are suitable for screening of overweight to prevent metabolic syndrome in elder Japanese workers. *J Occup Health*. 2003; 45(6): 335-343
6. Bernd Nilius, Joris Vriens, Jean Prenen, Guy Droogmans, and Thomas Voets. TRPV4 calcium entry channel: a paradigm for gating diversity. *Am J Physiol Cell Physiol*, 2004; 286: C195-C205
7. Caterina MJ, Schumacher MA, Tominaga M, et al. The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature*, 1997; 389: 816-824
8. Chung MK, Guler AD, Caterina MJ. Biphasic currents evoked by chemical or thermal activation of the heat-gated ion channel, TRPV3. *J Biol Chem*, 2005; 280(16): 15928-15941
9. Clapham DE, Runnels LW, Strübing C. The TRP ion channel family. *Nature Rev Neurosci*, 2001; 2:387-396
10. Gong DW, He Y, Karas M, et al. Uncoupling protein-3 is a mediator of thermogenesis regulated by thyroid hormone, beta3-adrenergic agonists, and leptin. *J Biol Chem*, 1997; 272(39): 24129-24132
11. Hu HZ, Gu Q, Wang C, Colton CK, Tang J, Kinoshita-Kawada M, Lee LY, Wood JD, Zhu MX. 2-aminoethoxydiphenyl borate is a common activator of TRPV1, TRPV2, and TRPV3. *J Biol Chem*, 2004; 279: 35741-35748
12. Montell C, Rubin GM. Molecular characterization of the drosophila trp locus: putative integral membrane protein required for phototransduction. *Neuron*, 1989; 2: 1313-1323
13. Muraki K, Iwata Y, Katanosaka Y, et al. TRPV2 is a component of osmotically sensitive cation channels in murine aortic myocytes. *Circ Res*, 2003; 93(9): 829-838
14. Sakurai T, Amemiya A, Ishii M, et al. Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell*, 1998; 92(5): 696-696
15. Schoeller DA, Van Santen E. Measurement of energy expenditure in humans by doubly labelled water method. *J Appl Physiol*, 1982; 53: 955-959
16. Simoneau JA, Kelley DE, Warden C. Obesity and increased contractile activity influence the protein content of UCP2 in human skeletal muscle. *Int J Obes Relat Metab Disord*, 1999; 23 Suppl 6: 68-71
17. Solanes G, Vidal-Puig A, Grujic D, et al. The human uncoupling protein-3 gene. Genomic structure, chromosomal localization, and genetic basis for short and long form transcripts. *J*

Biol Chem, 1997; 272(41): 25433-25436

18. Story GM, Peier AM, Reeve AJ, et al. ANKTM1, a TRP-like channel expressed in nociceptive neuron, is activated by cold temperature. *Cell*, 2003; 112: 819-829
19. Yu XX, Mao W, Zhong A, et al. Characterization of novel UCP5/BMCP1 isoforms and differential regulation of UCP4 and UCP5 expression through dietary or temperature manipulation. *FASEB J*, 2000; 14(11): 1611-1618
20. Zhang Y, Proenca R, Maffei M, et al. Positional cloning of the mouse obese gene and its human homologue. *Nature*, 1994; 372: 425-432